

SUPREP BOWEL PREP KIT

Date of procedure: ____ / ____ / ____

FOLLOW THIS INSTRUCTION SHEET, NOT THE INSTRUCTIONS ON THE BOX.

THE DAY BEFORE YOUR PROCEDURE...NO SOLID FOODS! CLEAR LIQUIDS ONLY!

On the day before your procedure...

What You CAN do:

- You may have clear liquids ONLY; please have nothing for dinner

What You CANNOT do:

- DO NOT** drink milk
- DO NOT** eat or drink anything colored red or purple
- DO NOT** drink alcoholic beverages

Any of the following liquids are OK to drink:

- Water
- Strained fruit juices (without pulp) including apple, orange, white grape, or white cranberry
- Limeade or lemonade
- Coffee or tea (**DO NOT** use any dairy or non-dairy creamer)
- Chicken broth
- Gelatin desserts without added fruit or topping (**NO RED OR PURPLE**)

Note:

- SUPREP Bowel Prep Kit is indicated for cleansing of the colon as a preparation for colonoscopy in adults
- Be sure to tell your doctor about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements. SUPREP Bowel Prep Kit may affect how other medicines work
- Medication taken by mouth may not be absorbed properly when taken within 1 hour before the start of each dose of SUPREP Bowel Prep Kit
- The most common adverse reactions after administration of SUPREP Bowel Prep Kit were overall discomfort, abdominal distention, abdominal pain, nausea, vomiting, and headache
- If you have any questions about taking SUPREP Bowel Prep Kit, call your doctor

To learn more about this product, please call 1-800-874-6756 or visit our Web site at www.suprepkit.com

Split-Dose (2-Day) Regimen

BEGIN STEP 1: BETWEEN 5-9PM

- On the evening before the procedure (or when your doctor tells you to begin) complete steps 1 through 4 using one (1) 6-ounce bottle before going to bed
- On the morning of your procedure, repeat steps 1 through 4 using the other 6-ounce bottle

STEP 1



Pour **ONE (1)** 6-ounce bottle of SUPREP liquid into the mixing container.

STEP 2



Add cool drinking water to the 16-ounce line on the container and mix.

STEP 3



Drink **ALL** the liquid in the container.

STEP 4



IMPORTANT
You **must** drink two (2) more 16-ounce containers of water over the next 1 hour.

MORNING OF: BEGIN STEP 1, 5 HRS BEFORE PROCEDURE

NOTE: Dilute the solution concentrate as directed prior to use.

NOTE: You **must** finish drinking the final glass of water at least 2 hours, or as directed, before your colonoscopy.

Provided as an educational service from **BrainTree**

©2015 BrainTree Laboratories, Inc.

SU-12253

December, 2015

Please read the Full Prescribing Information and Medication Guide in the kit.